




IDC 203: INTRODUCTION TO EARTH SCIENCES



**Table
2.4**

Physical Properties of Minerals

Property	Relation to Composition and Crystal Structure
Hardness	Strong chemical bonds give high hardness. Covalently bonded minerals are generally harder than ionically bonded minerals.
Cleavage	Cleavage is poor if bond strength in crystal structure is high and is good if bond strength is low. Covalent bonds generally give poor or no cleavage; ionic bonds are weak and so give excellent cleavage.
Fracture	Type is related to distribution of bond strengths across irregular surfaces other than cleavage planes.
Luster	Tends to be glassy for ionically bonded crystals, more variable for covalently bonded crystals.
Color	Determined by kinds of atoms and trace impurities. Many ionically bonded crystals are colorless. Iron tends to color strongly.
Streak	Color of fine powder is more characteristic than that of massive mineral because of uniformly small size of grains.
Density	Depends on atomic weight of atoms and their closeness of packing in crystal structure. Iron minerals and metals have high density; covalently bonded minerals have more open packing and so have lower density.

Outline

- Rocks
- Igneous rock- types and forms
- Tectonic settings of Igneous rock formation
- Sedimentary rock formation

Rocks

- A naturally occurring consolidated mixture of minerals or mineral-like substances
- Rocks can be also made up of non mineral matter (e.g., obsidian and coal)
- The identity of rock is determined partly by its **mineral** and partly by its **texture**.

Rocks

Constituent minerals

Orthoclase feldspar



Quartz




Biotite



Plagioclase feldspar



Rocks are naturally occurring aggregates of minerals.



Plagioclase feldspar
Orthoclase feldspar
Biotite
Quartz



Rock (granite)


Texture

- Texture describes size and shapes of a rocks mineral crystals or grains.
- The grain large enough to be seen the rock is classified as **coarse** grained.
- The grain **not** large enough to be seen the rock is classified as **fine** grained.

Rocks

- Rocks formed by solidification of molten rock are called **igneous rock**
- Weathering and erosion of rocks exposed at the surface – **sedimentary rocks**
- Rocks formed under high pressure and temperature in deep crust and upper mantle is **metamorphic rocks**

Rocks



Rocks

Table 3.5

Some Common Minerals
of Igneous, Sedimentary,
and Metamorphic Rocks

Igneous Rocks	Sedimentary Rocks	Metamorphic Rocks
*Quartz	*Quartz	*Quartz
*Feldspar	*Clay minerals	*Feldspar
*Mica	*Feldspar	*Mica
*Pyroxene	Calcite	*Garnet
*Amphibole	Dolomite	*Pyroxene
*Olivine	Gypsum	*Staurolite
	Halite	*Kyanite

Note: Asterisk indicates that the mineral is a silicate.

Igneous rocks

- **Formed from the cooling and consolidation of magma**
- **The first division of igneous rocks were made based on the evidences of texture**

Clues for early division of igneous rocks

- Volcanic rocks**
- Laboratory studies of crystallization**
- Granite as evidence of slow cooling**

Texture of igneous rocks

- **Controlled by cooling rate**
- **Grain size**
- **Degree of crystallinity**
- **Vesicularity**


Classification of igneous rocks

Defined by texture:

Fine-grained: extrusive or volcanic

Coarse-grained: intrusive or plutonic

Igneous Rocks



- Plutonic (intrusive) — cooled below the surface
- volcanic (extrusive) — cooled on the surface

Coarsely crystalline granite



Photomicrograph of granite




Fig. 4.1

Finely crystalline basalt




Fig. 4.1

Photomicrograph of basalt




Fig. 4.1

Igneous Classification by texture

Extrusive

Basalt

Andesite

Rhyolite

Intrusive

Gabbro

Diorite

Granite

Extrusive

Intrusive



Igneous textures

Glassy

no minerals present

Crystalline

rocks made of mineral grains

Porphyritic

mixture of coarse and fine


Vesicular

with bubble holes


Igneous textures



Glassy



Crystalline



Porphyritic



Vesicular

Classification of igneous rocks

- **Determined by composition (both chemical and mineralogical)**
- **Magnesium (Mg) + iron (Fe) = Mafic**
- **Feldspar + quartz (Si) = Felsic**

Classification of igneous rocks

Table 4.1

Common Minerals of Igneous Rocks

Compositional Group	Mineral	Chemical Composition	Silicate Structure
FELSIC	Quartz	SiO_2	Frameworks
	Potassium feldspar	KAlSi_3O_8	
	Plagioclase feldspar	$\left. \begin{array}{l} \text{NaAlSi}_3\text{O}_8 \\ \text{CaAl}_2\text{Si}_2\text{O}_8 \end{array} \right\}$	
	Muscovite (mica)	$\text{KAl}_3\text{Si}_3\text{O}_{10}(\text{OH})_2$	Sheets
MAFIC	Biotite (mica)	$\left. \begin{array}{l} \text{K} \\ \text{Mg} \\ \text{Fe} \\ \text{Al} \end{array} \right\} \text{Si}_3\text{O}_{10}(\text{OH})_2$	Double chains
	Amphibole group	$\left. \begin{array}{l} \text{Mg} \\ \text{Fe} \\ \text{Ca} \\ \text{Na} \end{array} \right\} \text{Si}_8\text{O}_{22}(\text{OH})_2$	
	Pyroxene group	$\left. \begin{array}{l} \text{Mg} \\ \text{Fe} \\ \text{Ca} \\ \text{Al} \end{array} \right\} \text{SiO}_3$	Single chains
	Olivine	$(\text{Mg,Fe})_2\text{SiO}_4$	Isolated tetrahedra


Classification of igneous rocks

When we talk about the chemical composition of a rock we usually speak in terms of the oxides, *e.g.*,


	Typical basalt	Typical granite
--	----------------	-----------------

SiO₂	50%	70%
Al₂O₃	15%	12%
FeO+MgO	15%	3%
CaO	8%	2%
K₂O+Na₂O	5%	8%

Classification of igneous rocks



Classification of igneous rocks



How do rocks melt?

Partial melting : the fraction of rock that has melted at given temperature

- **The temperature at the depth of crust and mantle**
- **Last minerals to form will melt at lowest temperature**
- **Biggest changes will be for small degrees of melting**

How do rocks melt?

- **The pressure due to overburden rocks also affect partial melting**
- **The melting temperature increases with high pressure**
- **Decompression melting: sudden decrease in pressure**

Increase in depth in crust and mantle raises temperature

Increase in pressure raises melting range temperatures

Increase in temperature raises proportion of partial melt

Increase in water content lowers melting temperature

Rock composition affects melting temperature (silicic, lower temperature mafic, higher temperature)

Factors affecting melting temperatures

The formation of magma chamber

- The large pools of molten rock form magma chamber
- The rise of magmas at rate from 0.3 m/year to almost 50 m/year

Magmatic differentiation

- The process by which rocks of varying chemical composition can arise from a uniform parent magma.
- This segregation happens in several way following a sequence described as **Bowen reaction series**


Fractional crystallization

- **The modification of magma by crystallization and removal of mineral phases.**
- **Because only certain elements will go into a given mineral, this will tend to change the composition of the remaining liquid.**

Bowen's reaction series

- **Series of chemical reactions that take place in silicate magmas as they cool**
- **First investigated in the 1920s and 1930s by N. L. Bowen**
- **Important experiments that help us understand the evolution of magmas**

Types of igneous structures



Forms of intrusive igneous masses

Plutons: large igneous bodies formed in deep earth crust

Plutons can be divided into two groups:

1) Concordant

2) Discordant

Forms of intrusive igneous masses

Concordant:

Sills

Discordant:

Dikes

Batholiths

Forms of intrusive igneous masses

Batholith: Any deep-seated pluton of coarse-grained rocks that has a surface exposure of over 100 sq. km that is mostly granitic. Examples include Sierra Nevada, Coast Range, Idaho batholiths.


Stock: Same as a batholith, only smaller

Forms of intrusive igneous masses

Sills: sheetlike body formed by the injection of magma between parallel layers of bedded country rock.

Dyke: sheetlike body formed by the injection of magma cut across the layers in bedded country rock.

Sill



Dike



Forms of intrusive igneous masses


Veins: deposits of minerals found within a rock fracture that are foreign to the country rock.

Veins



Monochromatic chert cut by quartz veins

Pegmatite veins



Fig

Igneous processes and plate tectonics

Sedimentary rocks

Weathering –Decomposition and disintegration of pre-existing rock into small fragments or new minerals

Transportation of the sediments to a sedimentary basin

Deposition of the sediment

Burial and Lithification to make sedimentary rock

Sedimentary stages in the rock cycle




Fig. 7.1

Sedimentary rocks

Weathering process

Types of sedimentary rocks

Depositional environments

Weathering process

1. Physical (Mechanical): Large rocks broken into smaller fragments with **no change in composition**
2. Chemical: Rocks dissolved –
chemical and mineralogical composition can be altered
 - new minerals may form
3. Biological : plants & animals


Physical weathering

1. **Joints** (also termed extensional fractures) are planes of separation on which no shear displacement has taken place



Physical Weathering

2. Frost wedging



The black lines in the rock represent fractures that are occurring in the rock.

The blue lines in the rock represent water soaking into the fractures.

The water freezes and expands. If this cycle of freezing, expansion, and thawing continues, the rock will gradually disintegrate.

Physical Weathering

3. Salt precipitation



Physical Weathering

4. Abrasion

Physical grinding of rock fragments



Chemical Weathering


Chemical weathering is the weakening and subsequent disintegration of rock by chemical reactions.

These reactions include

- i) Hydration
- ii) Hydrolysis
- iii) Carbonation

Chemical Weathering

Carbonation is the process of rock minerals reacting with carbonic acid. Carbonic acid is formed when water combines with carbon dioxide. Carbonic acid dissolves or breaks down minerals in the rock.



(carbon dioxide + water → carbonic acid)



(calcite + carbonic acid → calcium + bicarbonate)

Chemical Weathering

Hydrolysis is a chemical reaction caused by water.

Water changes the **chemical composition and size of minerals in rock**, making them less resistant to weathering.



Chemical Weathering

Hydration is the absorption of water into the mineral structure.

A good example of hydration is the absorption of water by anhydrite, resulting in the formation of gypsum. Hydration expands volume and also results in rock deformation.

Biological Weathering


- This form of weathering is caused by activities of living organism



Factors affect the rates of weathering process

- **Climate**
- **Structural weakness**
- **Topography**
- **Time**
- **Nature of rocks and minerals**

Weathering process



**Table
7.1**

Minerals Remaining in Clastic Sediments Derived from an Average Granite Outcrop Under Varying Intensities of Weathering

Intensity of Weathering		
Low	Medium	High
Quartz	Quartz	Quartz
Feldspar	Feldspar	Clay minerals
Mica	Mica	
Pyroxene	Clay minerals	
Amphibole		


HOW COMMON ARE SEDIMENTARY ROCKS?





Crustal volume



Land surface area



 Igneous

 Sedimentary

 Metamorphic

Sedimentary Rocks

- *Detrital (clastic) rocks* produced from rock fragments
- *Chemical rocks* produced by precipitation of dissolved ions in water
- *Organic rocks* produced by accumulation of biological debris, such as in swamps or bogs

Sedimentary rock types and *sedimentary structures* within the rocks give clues to *past environments*

Transport and deposition of Clastic sediments

- **Movement of sediment by wind, ice or water.**
- **Mode of transport produces distinctive deposits.**


Transport affects the sediment in several ways

Sorting: measure of the variation in the range of grain sizes in a clastic rock or sediment


- **Well-sorted sediments indicate that they have been subjected to prolonged water or wind action.**
- **Poorly-sorted sediments are either not far-removed from their source or deposited by glaciers.**

Sorting

Well-sorted



Poorly-sorted



Well-sorted sand



Poorly-sorted sand




Transport affects the sediment in several ways

Roundness: measure of how rounded the corners are

Sphericity: measure of how much it is like a sphere

Sorting, roundness, and sphericity all increase with amount of transport.

Roundness and sphericity



Types of detrital rocks

Largely based on the size of the particles, which may be anything.

- *Sediment* -classified by *particle size*
 - Boulder - >256 mm
 - Cobble - 64 to 256 mm
 - Pebble - 2 to 64 mm ----- Gravel
 - Sand - 1/16 to 2 mm
 - Silt - 1/256 to 1/16 mm
 - Clay - <1/256 mm

Types of detrital rocks

Conglomerate

Breccia

Sandstone **(quartzite, arkose, greywacke)**

Shale

Mudstone

Siltstone

Clastic sedimentary rocks

- *Breccia and Conglomerate*
 - *Coarse-grained clastic* sedimentary rocks
 - Sedimentary breccia composed of coarse, *angular rock fragments* cemented together
 - Conglomerate composed of *rounded gravel* cemented together

Conglomerate



Breccia



Clastic sedimentary rocks

- *Sandstone*

- *Medium-grained clastic* sedimentary rock

- Types determined by composition

- *Quartz sandstone* - >90% quartz grains


- *Arkose* - mostly feldspar and quartz grains

- *Graywacke* - sand grains surrounded by dark, fine-grained matrix, often clay-rich


Sandstone



Breck Kent



(a) Quartz arenite:
pure quartz



(b) Arkose:
feldspar-rich



(c) Lithic sandstone:
rock-fragment-rich



(d) Graywacke:
matrix-rich



Four major groups of sandstones

Shale

Shales are clastic rocks, made up mainly fine silt/clay

They are most abundant sedimentary rocks, accounts for about 80% of them

Often contain fossils

Mostly hydrous aluminum silicate in composition = from weathered feldspars

Deposition takes place under low fluvial regime or under weak water current.

Eg. Offshore or in Lagoon

- Shales are made of fine well sorted silt and clayey sediments, where normally one can expect high porosity and permeability.

Shale



Types of chemical sedimentary rocks

Limestone



Chert



Salt



Gypsum



Coal

altered organic debris

Chemical sedimentary rocks

- *Carbonates*
 - Contain CO_3 as part of their chemical composition
 - *Limestone* is composed mainly of *calcite*
 - Most are *biochemical*, but can be *inorganic*
 - Often contain easily recognizable fossils

Chemical sedimentary rocks

Limestone: It is a non-clastic rock formed either chemically or due to precipitation of calcite (CaCO_3) from organisms usually shell. These remains will result in formation of a limestone.


Limestones formed by chemical precipitation are usually fine grained, whereas, in case of organic limestone the grain size vary depending upon the type of organism responsible for the formation

Fossiliferous Limestone: which medium to coarse grained, as it is formed out of cementation of Shells.


Limestone



Foraminifera in the eye of a needle



Fossiliferous limestone



Chemical sedimentary rocks

Chert

- Hard, compact, fine-grained, formed almost entirely of silica
- Can occur as layers or as lumpy nodules within other sedimentary rocks, especially limestones



Evaporites

- These rocks are formed within the a depositional basin from chemical substances dissolved in the seawater or lake water
- Characteristic of arid conditions

Evaporites

Minerals precipitate according to solubility.

Gypsum

50%



$\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$


Halite

90%



NaCl


Relative abundance of sedimentary rock types



Sedimentary environment

- Sediments accumulate in some environment of deposition or **depositional environments**
- These areas receive net deposition
- Erosion may occur, but deposition dominates
- Features of these depositional environments are preserved in the rock record
- Examples:
 - Sediment texture
 - Fossils of organisms that lived in the environment
- Ancient environments can be reconstructed from the clues that are preserved in the sedimentary rocks

Common sedimentary environments



GEOGRAPHIC LOCATION:

Shoreline

TRANSPORT AGENT:

Waves and tides

TYPE OF MEDIUM:

Seawater

DEPTH OF WATER:

0–5 meters

**ORGANISMS THAT
MODIFY SEDIMENT:**

Burrowing invertebrates

CLIMATE:

Tropical

PLATE-TECTONIC SETTING:

Plate convergence zone

S
E
D
I
M
E
N
T
A
R
Y
E
N
V
I
R
O
N
M
E
N
T

B
e
a
c
h

SEDIMENT
DEPOSITED:

Sand and gravel

Characteristics of a sedimentary environment

**Table
7.2**

Clastic Sedimentary Environments

Environment	Agent of Transportation, Deposition	Sediments
CONTINENTAL		
Alluvial	Rivers	Sand, gravel, mud
Desert	Wind	Sand, dust
Lake	Lake currents, waves	Sand, mud
Glacial	Ice	Sand, gravel, mud
SHORELINE		
Delta	River + waves, tides	Sand, mud
Beach	Waves, tides	Sand, gravel
Tidal flats	Currents	Sand, mud
MARINE		
Continental shelf	Waves, tides	Sand, mud
Continental margin	Ocean currents	Mud, sand
Deep sea	Ocean currents, settling	Mud

**Table
7.3**

Major Chemical and Biochemical Sedimentary Environments

Environment	Agent of Precipitation	Sediments
SHORELINE AND MARINE Carbonate (includes reef, bank, deep sea, etc.)	Shelled organisms, some algae; inorganic precipitation from seawater	Carbonate sands and muds, reefs
Evaporite	Evaporation of seawater	Gypsum, halite, other salts
Siliceous: deep sea	Shelled organisms	Silica
CONTINENTAL Evaporite	Evaporation of lake water	Halite, borates, nitrates, other salts
Swamp	Vegetation	Peat