Sets

Dhruva Sambrani

20 January, 2020

Sets

Set is a well defined collection of objects. Well defined here means that an element of the Domain x should be unambiguously belong to or not belong to set S.

 $\mathbf{x} \in \mathbf{A} \equiv$ "x belongs to set A"

 $x \notin A \equiv$ "x does not belong to set A"

Properties

- All elements unique.
- Unordered.

Types

Sets need not be of a particular "type" - {{1,2,3},6,7} May also be infinite. {1,2,3,4....} Null/Empty set = {} = \emptyset

Operations on sets

- Subset: $A \subseteq B$. "A is a subset of B." $x \in A \Longrightarrow x \in B$.
- Equality: A=B iff A⊆B \land B⊆A. "A is equal to B." x∈A \implies x∈B \land x∉A \implies x∉B
- Proper Subset: $A \subset B \ x \in A \implies x \in B \land A \neq B$.
- Complement: U\A OR Ac. Exactly all x's in U NOT in A.
- Union: $A \cup B$. "A union B". Exactly all elements in $A \vee B$.
 - If $\{A\alpha\}$ is a collection of sets indexed by I, then $\bigcup A\alpha = \text{set of } x \text{ st } x \in \alpha 0$ for some $\alpha \in I$.
- Intersection: A∩B "A intersection B". Contains exactly all elements in A AND B.

– x
є $\bigcap\!A\alpha$ iff x $\forall\alpha{\in}I,$ x
<A\alpha

- Cartesian Product: $A \times B$ "Cartesian Product of A and B". Its the collection of all 2 element sequences (a,b) st $a \in A$, $b \in B$
 - A1×A2×...×An is the collection of all n element sequences (ai) st ai
∈Ai
- Relation operator: aRb "a is related to b". A relation is a subset of A×B.
- Function: f: A \rightarrow B. It is a relation such that
 - 1. $\forall a \in A, \exists b \text{ st aRb.}$
 - 2. aRb and aRc \Longrightarrow b=c.
 - Equality of functions f=g \implies f⊆g \land g⊆f.

Languages

 Σ is a finite set called an "alphabet".

The finite sequence is called a string.

A language over Σ is a set of strings.