

**IISER Mohali**  
**MTH102: Analysis in One Variable**  
**Homework No. 06**  
**To be discussed during tutorial on 04 March, 2016**

- Please solve all the problems.
- Tutors will discuss **tutorial problems** during tutorial sessions.
- If time permits, tutors may also discuss **extra problems** during tutorial sessions.

**Tutorial Problems:**

- (1) Use the  $\epsilon - \delta$  definition of uniform continuity to show that the function  $f : [0, 2] \rightarrow \mathbb{R}$  defined by  $f(x) = x^2$  is uniformly continuous.
- (2) Show that the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(x) = x^3$  is not uniformly continuous.  
Remark: In fact, the only polynomials that are uniformly continuous on  $\mathbb{R}$  are those of degree at most 1.
- (3) For each of the following power series, determine the radius of convergence and also the exact interval of convergence:
  - (a)  $\sum_{n=0}^{\infty} \sqrt{n}x^n$
  - (b)  $\sum_{n=0}^{\infty} \frac{2^n}{n^2}x^n$
- (4) Let  $\sum_{n=0}^{\infty} a_n x^n$  be a power series with radius of convergence  $R$ . Suppose that all the coefficients  $a_n$  are integers and all but finitely many  $a_n$ 's are non-zero. Then prove that  $R \leq 1$ .  
Hint: Use the definition of radius of convergence.
- (5) Give an example of a power series whose exact interval of convergence is  $(-1, 1]$ .  
Hint: Take a power series which becomes harmonic series at 1 or -1.
- (6) Let  $\sum_{n=0}^{\infty} a_n x^n$  and  $\sum_{n=0}^{\infty} b_n x^n$  be two power series with radius of convergence  $R_1$  and  $R_2$ , respectively. Define their sum as

$$\sum_{n=0}^{\infty} a_n x^n + \sum_{n=0}^{\infty} b_n x^n = \sum_{n=0}^{\infty} (a_n + b_n) x^n.$$

- Prove that if  $R$  is the radius of convergence of  $\sum_{n=0}^{\infty} (a_n + b_n) x^n$ , then  $R \geq \min\{R_1, R_2\}$ .
- (7) In problem 5 above, give examples of power series where  $R = \min\{R_1, R_2\}$ .

**Extra Problems:**

- (1) Use the  $\epsilon - \delta$  definition of uniform continuity to show that the function  $f : [0, 2] \rightarrow \mathbb{R}$  defined by  $f(x) = \frac{x}{x+2}$  is uniformly continuous on  $[0, 2]$ .
- (2) Let  $f : S \rightarrow \mathbb{R}$  be a uniformly continuous function on a subset  $S$  of  $\mathbb{R}$ . Show that if  $(x_n)$  is a Cauchy sequence in  $S$ , then  $(f(x_n))$  is also a Cauchy sequence in  $\mathbb{R}$ .
- (3) For each of the following power series, determine the radius of convergence and also the exact interval of convergence:
  - (a)  $\sum_{n=0}^{\infty} n^2 x^n$
  - (b)  $\sum_{n=0}^{\infty} \left(\frac{x}{n}\right)^n$
- (4) In problem 5 above, give examples of power series where  $R > \min\{R_1, R_2\}$ .