



Differential Geometry

MTH201



EDITED BY
ADITYA DEV



Contents

1	Curves in the plane and in space	2
1.1	What is a curve?	2
1.2	Arc-Length	2
1.3	Reparametrization of a curve	3
1.4	Closed Curves	4
2	Curvature	6
2.1	What is curvature?	6
2.2	2D and 3D curves	6
2.2.1	Plane curve	6
2.2.2	Space curve	8
3	Global properties of curves	11
3.1	Simple closed curves	11
3.2	The isoperimetric inequality	11
3.3	The four vertex Theorem	12
4	3D Surfaces	13
4.1	What is a surface?	13
4.2	Smooth Surfaces	13
4.3	Smooth Map	14
4.4	Tangents and derivatives	15

Chapter 1

Curves in the plane and in space

§1.1 What is a curve?

Definition 1.1.1: Parametrized curve

A *parametrized curve* in \mathbb{R}^n is a map $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^n$, for some $(\alpha, \beta) \subseteq \mathbb{R}$

Example : $\gamma(t) : (-\infty, \infty) \rightarrow (t, t^2)$

Note There can be different parametrizations for the same curve; but it's not mandatory that they have same properties.

Smooth Function A function $f : (\alpha, \beta) \rightarrow \mathbb{R}$ is said to be smooth if the derivative $\frac{d^n f}{dt^n}$ exists for all $n \geq 1$ and all $t \in (\alpha, \beta)$.

Definition 1.1.2: Tangent Vector

If γ is a parametrized curve, its first derivative $\dot{\gamma}(t)$ is called the tangent vector of γ at the point $\gamma(t)$.

Proposition 1.1.1

If the tangent vector of a parametrized curve is constant, the image of the curve is (part of) a straight line.

§1.2 Arc-Length

Recall that if $\mathbf{v} = (v_1, v_2, \dots, v_n) \in \mathbb{R}^n$, then it's length is:

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$$

If \mathbf{u} is another vector in \mathbb{R}^n , $\|\mathbf{u} - \mathbf{v}\|$ is the length of the straight line segment joining the points \mathbf{u} and \mathbf{v} in \mathbb{R}^n .

Definition 1.2.1: Arc-length

The *arc-length* of a curve γ starting at the point $\gamma(t_0)$ is the function $s(t)$ given

by:

$$s(t) = \int_{t_0}^t \|\dot{\gamma}(x)\| dx$$

Note that if we choose a different starting point, then the new arc-length differs from the previous one (*but how much?*)

Definition 1.2.2

If $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ is a parametrized curve, its speed at point $\gamma(t)$ is $\|\dot{\gamma}(t)\|$, and γ is said to be a unit-speed curve if $\dot{\gamma}(t)$ is a unit-vector $\forall t \in (\alpha, \beta)$.

Proposition 1.2.1

Let $\mathbf{n}(t)$ be a unit vector that is a smooth function of a parameter t . Then, the dot product

$$\mathbf{n}(t) \cdot \dot{\mathbf{n}}(t) = 0 \quad \forall t$$

so, either $\dot{\mathbf{n}}(t)$ is zero or perpendicular to $\mathbf{n}(t)$

§1.3 Reparametrization of a curve

Definition 1.3.1: Reparametrization

A parametrized curve $\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^n$ is a reparametrization of a parametrized curve $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^n$ if \exists a smooth bijective map $\phi : (\tilde{\alpha}, \tilde{\beta}) \rightarrow (\alpha, \beta)$ (the reparametrization map) such that the inverse map $\phi^{-1} : (\alpha, \beta) \rightarrow (\tilde{\alpha}, \tilde{\beta})$ is also smooth and

$$\tilde{\gamma}(\tilde{t}) = \gamma(\phi(\tilde{t}))$$

Definition 1.3.2: Regular Curve

A point $\gamma(t)$ of a parametrized curve γ is called a regular point if $\dot{\gamma}(t) \neq 0$ otherwise $\gamma(t)$ is a singular point of γ . A curve is regular if all of its points are regular

Proposition 1.3.1

Any reparametrization of a regular curve is regular.

Proposition 1.3.2

If $\gamma(t)$ is a regular curve, its arc-length s , starting at any point of γ , is smooth function of t .

Theorem 1.3.1: Unit-speed reparametrization

A parametrized curve has a unit-speed reparametrization if and only if it is regular.

Corollary 1.3.1

Let γ be a regular curve and let $\tilde{\gamma}$ be a unit-speed reparametrization of γ :

$$\tilde{\gamma}(u(t)) = \gamma(t) \quad \forall t$$

where u is a smooth function of t . Then, if s is the arc-length of γ (starting at any point), we have:

$$u = \pm s + c \quad \text{for some } c \in \mathbb{R} \quad (1.1)$$

Conversely, if u is given by Eq. 1.1 for some value of c and with either sign, then $\tilde{\gamma}$ is a unit-speed reparametrization of γ .

§1.4 Closed Curves

Definition 1.4.1: Periodic Curve

Let $\gamma : \mathbb{R} \rightarrow \mathbb{R}^n$ be a smooth curve and let $T \in \mathbb{R}$. We say that γ is T -periodic if:

$$\gamma(t + T) = \gamma(t) \quad \forall t \in \mathbb{R}$$

If γ is not constant and is T -periodic for some $T \neq 0$, then γ is said to be closed.

Note if γ is T -periodic then it is $-T$ -periodic too because

$$\gamma(t - T) = \gamma(t - T + T) = \gamma(t)$$

It follows that if γ is T -periodic for some $T \neq 0$, then it is T -periodic for some ($T > 0$).

Definition 1.4.2: Self-intersection

A curve γ is said to have a self-intersection at a point \mathbf{p} of the curve if there exist parameter values $a \neq b$ such that

- $\gamma(a) = \gamma(b) = \mathbf{p}$
- if γ is closed with period T , then $a - b$ is not an integer multiple of T .

Proof. Assume there exists no lower bound for the period of curve. Then if T_1 is the period of γ then $\exists T_2$ such that T_2 is also the period of the curve and by iteration we can show:

$$T_1 > T_2 > T_3 \dots > 0$$

is a sequence of the periods for curve γ

Since, the sequence is bounded and monotonic, therefore $\lim_{r \rightarrow \infty} T_r = T$ i.e the sequence is convergent \implies the sequence is cauchy.

And by definition of cauchy sequence:

$$\forall \epsilon > 0 \exists N : \forall m, n > N \implies |T_m - T_n| < \epsilon$$

Let, $T_m > T_n$ (won't change the definition). Also, we know that if T_m & T_n are periods of $\gamma \implies T_m - T_n \implies T_m = T_n + \epsilon$ is also the period of gamma (trivial to prove!).

$$\gamma(t + T_n + \epsilon) = \gamma(t)$$

$$\gamma(t + \epsilon) = \gamma(t)$$

Since, it's true $\forall \epsilon > 0 \implies \gamma$ is constant. Hence, a contradiction (γ is non-constant.). So, our assumption was false. ■

Chapter 2

Curvature

§2.1 What is curvature?

Definition 2.1.1: Curvature

If γ is a unit-speed curve with parameter t , its curvature $\kappa(t)$ at the point $\gamma(t)$ is defined to be $\|\ddot{\gamma}(t)\|$.

Note we have defined curvature for a unit-speed parametric only.

Theorem 2.1.1

The curvature for any regular curve γ is given as

$$\kappa = \frac{\|(\dot{\gamma} \cdot \dot{\gamma})\ddot{\gamma} - (\dot{\gamma} \cdot \ddot{\gamma})\dot{\gamma}\|}{\|\dot{\gamma}\|^4}$$

Proposition 2.1.1

Let $\gamma(t)$ be a regular curve in \mathbb{R}^3 . Then its curvature is

$$\kappa = \frac{\|\ddot{\gamma} \times \dot{\gamma}\|}{\|\dot{\gamma}\|^3}$$

where \times is our usual vector cross product.

Problem 1. Show that, if the curvature $\kappa(t)$ of a regular curve $\gamma(t)$ is > 0 every- where, then $\kappa(t)$ is a smooth function of t . Give an example to show that this may not be the case without the assumption that $\kappa(t) > 0$.

§2.2 2D and 3D curves

§2.2.1 Plane curve

Let γ be a unit-speed curve in a plane. And let the tangent vector be

$$\mathbf{t} = \dot{\gamma}$$

Note, \mathbf{t} is a unit-vector. There are two vectors perpendicular to \mathbf{t} ; we make a choice by defining \mathbf{n}_s , the *signed unit normal* of γ , to be the unit vector obtained by rotating \mathbf{t} anti-clockwise by $\pi/2$

So, by Proposition 1.2.1, $\dot{\mathbf{t}} = \dot{\gamma}$ is perpendicular to \mathbf{t} , and hence parallel to \mathbf{n}_s . Thus, there is a scalar κ_s such that

$$\dot{\gamma} = \kappa_s \mathbf{n}_s$$

κ_s is called the signed curvature of γ . And since $\|\mathbf{n}_s\| = 1$, we have

$$\kappa = \|\kappa_s \mathbf{n}_s\| = |\kappa_s|$$

Note, we have defined the signed curvature for unit-speed curve. If γ is any regular curve, then we define the above defined parameters to be those of its unit speed parametrization.

Intuitively, since $\gamma(t)$ is assumed to be a unit-speed curve on a plane, then $\dot{\gamma}(t)$ can be measured by angle $\phi(s)$ such that:

$$\dot{\gamma}(s) = (\cos \phi(s), \sin \phi(s)) \quad (2)$$

Also, we can think curvature as rate at which the angle of tangent vector is changing, so if we find the derivative of the above curve then it simply represents a changing angle parameter.

Proposition 2.2.1

Let $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ be a unit speed curve, let $s_0 \in (\alpha, \beta)$ and let ϕ_0 be such that

$$\dot{\gamma}(s_0) = (\cos \phi_0, \sin \phi_0)$$

Then $\exists!$ smooth function: $\phi : (\alpha, \beta) \rightarrow \mathbb{R}$ such that $\phi(s_0) = \phi_0$ and that Eq. (2) holds for all $s \in (\alpha, \beta)$

Definition 2.2.1: Turning Angle

The smooth function ϕ in Proposition 2.2.1 is called the turning angle of γ determined by the condition $\phi(s_0) = \phi_0$.

Proposition 2.2.2

Let $\gamma(s)$ be a unit-speed plane curve, and let $\phi(s)$ be a turning angle for γ . Then,

$$\kappa_s = \frac{d\phi}{ds}$$

Thus, **the signed curvature is the rate at which the tangent vector of the curve rotates.**

Corollary 2.2.1

The total signed curvature of a closed plane curve is an integer multiple of 2π

The next result shows that a unit-speed plane curve is essentially determined once we know its signed curvature at each point of the curve. The meaning of ‘essentially’ here is ‘up to a direct isometry of \mathbb{R}^2 ’, i.e., a map $M : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ of the form

$$M = T_a \circ \rho_\theta$$

where ρ_θ is an anti-clockwise rotation by angle θ about the origin,

$$\rho_\theta = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta)$$

and T_a is a translation by vector \mathbf{a}

$$T_a(\mathbf{v}) = \mathbf{v} + \mathbf{a}$$

for any vector (x, y) and \mathbf{v} in \mathbb{R}^2

Theorem 2.2.1

Let $k : (\alpha, \beta) \rightarrow \mathbb{R}$ be any smooth function. Then, there is a unit-speed curve $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ whose signed curvature is k .

Further, if $\tilde{\gamma} : (\alpha, \beta) \rightarrow \mathbb{R}^2$ is any other unit-speed curve whose signed curvature is k , there is a direct isometry M of \mathbb{R}^2 such that

$$\tilde{\gamma}(s) = M(\gamma(s)) \quad \forall s \in (\alpha, \beta)$$

§2.2.2 Space curve

Definition 2.2.2

Let $\gamma(s)$ be a unit-speed curve in \mathbb{R}^3 , and let $\mathbf{t} = \dot{\gamma}$ be its unit tangent vector. If the curvature κ_s is non-zero, we define the principal normal of γ at the point $\gamma(s)$ to be the vector

$$\mathbf{n}(s) = \frac{1}{\kappa(s)} \dot{\mathbf{t}}(s)$$

Further, since $\|\dot{\mathbf{t}}\| = \kappa$, so \mathbf{n} is a unit-vector. Therefore by Proposition 1.2.1, so \mathbf{t} and \mathbf{n} are perpendicular. So

$$\mathbf{b} = \mathbf{t} \times \mathbf{n}$$

is a unit-vector perpendicular to both \mathbf{t} and \mathbf{n} . The vector $\mathbf{b}(s)$ is called the *binormal* vector of γ at point $\gamma(s)$. Thus, $\{\mathbf{t}, \mathbf{n}, \mathbf{b}\}$ is an orthonormal basis of \mathbb{R}^3 , and is *right-handed*.

Definition 2.2.3

From Definition 2.2.2 we have

$$\mathbf{b} = \mathbf{t} \times \mathbf{n}$$

Differentiating both sides gives

$$\dot{\mathbf{b}} = \dot{\mathbf{t}} \times \mathbf{n} + \mathbf{t} \times \dot{\mathbf{n}} = \mathbf{t} \times \dot{\mathbf{n}} \quad (3)$$

Equation (3) shows that $\dot{\mathbf{b}}$ is perpendicular to \mathbf{t} . Being perpendicular to both \mathbf{t}

and \mathbf{b} , $\dot{\mathbf{b}}$ must be parallel to \mathbf{n} , so

$$\dot{\mathbf{b}} = -\tau \times \mathbf{n}$$

for some scalar τ , which is called the *torsion* of γ .

Note that the torsion is only defined if the curvature is non-zero.

Definition 2.2.4

Let $\gamma(t)$ be a regular curve in \mathbb{R}^3 with nowhere-vanishing curvature. Then, denoting $\frac{d}{dt}$ by a dot, its torsion is given by

$$\tau = \frac{(\dot{\gamma} \times \ddot{\gamma}) \cdot \ddot{\gamma}}{\|\dot{\gamma} \times \ddot{\gamma}\|^2}$$

Proposition 2.2.3

Let γ be a regular curve in \mathbb{R}^3 with nowhere vanishing curvature (so that the torsion τ of γ is defined). Then, the image of γ is contained in a plane if and only if τ is zero at every point of the curve.

Theorem 2.2.2

Let γ be a unit-speed curve in \mathbb{R}^3 with nowhere vanishing curvature. Then,

$$\begin{aligned}\dot{\mathbf{t}} &= \kappa \mathbf{n} \\ \dot{\mathbf{n}} &= -\kappa \mathbf{t} + \tau \mathbf{b} \\ \dot{\mathbf{b}} &= -\tau \mathbf{n}\end{aligned}$$

The above equation is called as *Frenet-Serret equations*. Notice that the matrix

$$\begin{pmatrix} 0 & \kappa & 0 \\ -\kappa & 0 & \tau \\ 0 & -\tau & 0 \end{pmatrix}$$

which is the matrix of linear transformation is a *skew-symmetric*.

Proposition 2.2.4

Let γ be a unit-speed curve in \mathbb{R}^3 with constant curvature and zero torsion. Then, γ is a parametrization of (part of) a circle.

Theorem 2.2.3

Let $\gamma(s)$ and $\tilde{\gamma}(s)$ be two unit-speed curves in \mathbb{R}^3 with the same curvature $\kappa(s) > 0$ and the same torsion $\tau(s)$ for all s . Then, there is a direct isometry M of \mathbb{R}^3 such

that

$$\tilde{\gamma}(s) = M(\gamma(s)) \quad \forall s$$

Further, if k and t are smooth functions with $k > 0$ everywhere, there is a unit-speed curve in \mathbb{R}^3 whose curvature is k and whose torsion is t .

Chapter 3

Global properties of curves

§3.1 Simple closed curves

Definition 3.1.1

A *simple closed curve* in \mathbb{R}^2 is a closed curve in \mathbb{R}^2 that has no self-intersections.

Theorem 3.1.1: Jordan Curve Theorem

The complement of the image of γ (i.e., the set of points of \mathbb{R}^2 that are not in the image of γ) is the disjoint union of two subsets of \mathbb{R}^2 , denoted by $\text{int}(\gamma)$ and $\text{ext}(\gamma)$, with the following properties:

- $\text{int}(\gamma)$ is bounded, i.e. it is contained in the circle of sufficiently large radius.
- $\text{ext}(\gamma)$ is unbounded.
- Both the regions $\text{int}(\gamma)$ and $\text{ext}(\gamma)$ are connected, i.e they have the property that any two points in the same region can be joined by a curve contained entirely in the region.

Theorem 3.1.2: Hopf's Umlaufsatz

The total signed curvature of a simple closed curve in \mathbb{R}^2 is $\pm 2\pi$.

§3.2 The isoperimetric inequality

Definition 3.2.1: Area of a curve

The area contained by a simple closed curve γ is

$$\mathcal{A}(\gamma) = \int_{\text{int}(\gamma)} dx dy$$

Theorem 3.2.1: Green's Theorem

Let $f(x, y)$ and $g(x, y)$ be smooth functions (i.e., functions with continuous partial derivatives of all orders), and let γ be a positively-oriented simple closed curve.

Then,

$$\int_{\text{int}(\gamma)} \left(\frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \right) dx dy = \int_{\gamma} f(x, y) dx + g(x, y) dy$$

Proposition 3.2.1

If $\gamma(t) = (x(t), y(t))$ is a positively-oriented simple closed curve in \mathbb{R}^2 with period T , then

$$\mathcal{A}(\gamma) = \frac{1}{2} \int_0^T (x\dot{y} - y\dot{x}) dt$$

Theorem 3.2.2: Isoperimetric Inequality

Let γ be a simple closed curve, let $l(\gamma)$ be its length and let $\mathcal{A}(\gamma)$ be the area contained by it. Then,

$$\mathcal{A}(\gamma) \leq \frac{1}{4\pi} l(\gamma)^2$$

and equality holds if and only if γ is a circle.

§3.3 The four vertex Theorem

Definition 3.3.1: Vertex

A *vertex* of a curve $\gamma(t)$ in \mathbb{R}^2 is a point where its signed curvature κ_s has a stationary point, i.e., where $\frac{d\kappa_s}{dt} = 0$.

Theorem 3.3.1: Four Vertex Theorem

Every convex simple closed curve in \mathbb{R}^2 has at least four vertices.

Chapter 4

3D Surfaces

§4.1 What is a surface?

As in the case of curves, we make two definitions of the concept of surface. One of them (regular surface) emphasizes the fact that a surface, as we think of it, is a set of points. The other (parametrised surface) emphasizes the parametrization of the surface. While these two concepts were similar in the case of curves (every regular curve can be covered with a single parametrization, so it is a parametrised regular curve), they are different for surfaces: a sphere, for example, is a regular surface, but not a parametrised regular surface. We will further show that we need two parametric maps to describe the whole surface of a sphere and to keep it consistent with the other properties such as tangents and normals etc.

Definition 4.1.1: Surface

A subset S of \mathbb{R}^3 is a surface if, for every point $\mathbf{p} \in S$, there is an open set $U \subseteq \mathbb{R}^2$ and an open set $W \subseteq \mathbb{R}^3$ containing \mathbf{p} such that $S \cap W$ is homeomorphic to U i.e.

$$\sigma : U \subseteq \mathbb{R}^2 \rightarrow S \cap W$$

such that $\exists (u, v) \in U : \sigma(u, v) = \mathbf{p}$.

- 1 A subset of S of the form $S \cap W$, where W is an open subset of \mathbb{R}^3 , is called an open subset of S .
- 2 A continuous bijective function between two topological space (i.e. shapes here) is termed as *homeomorphism*

Definition 4.1.2: Surface Patch

A homeomorphism $\sigma : U \rightarrow S \cap W$ as in previous definition is called a *surface patch or parametrization* of the open subset $S \cap W$ of S .

A surface is some subset of \mathbb{R}^3 that can be covered by surface patches. Each surface patch looks like a (maybe deformed) piece of \mathbb{R}^2 .

§4.2 Smooth Surfaces

Smooth Functions If U is an open subset of \mathbb{R}^m , we say that a map $f : U \rightarrow \mathbb{R}^n$ is smooth if each of the n components of f , which are functions $U \rightarrow \mathbb{R}$, have continuous partial derivatives of all orders.

Definition 4.2.1: Regular Surface Patch

A surface patch $\sigma : U \rightarrow \mathbb{R}^3$ is called regular if it is smooth and the vectors σ_u and σ_v are linearly independent at all points $(u, v) \in \mathbb{R}^2$. Equivalently, σ should be smooth and the vector product $\sigma_u \times \sigma_v$ should be non-zero at every point of U .

Definition 4.2.2: Allowable Patch

If S is a surface, an allowable surface patch for S is a regular surface patch $\sigma : U \rightarrow \mathbb{R}^3$ such that σ is a homeomorphism from U to an open subset of S .

Definition 4.2.3: Smooth Surface

A smooth surface is a surface S such that, for any point $\mathbf{p} \in S$ there is an allowable surface patch σ as above such that $\mathbf{p} \in \sigma(U)$.

Definition 4.2.4: Atlas

A collection \mathcal{A} of allowable surface patches for a surface S such that every point of S is in the image of at least one patch in \mathcal{A} is called an atlas for the smooth surface S .

Proposition 4.2.1

The transition maps of a smooth surface are smooth.

Proposition 4.2.2

Let U and \tilde{U} be open subsets of \mathbb{R}^2 and let $\sigma : U \rightarrow \mathbb{R}^3$ be a regular surface patch. Let $\Phi : \tilde{U} \rightarrow U$ be a bijective smooth map with smooth inverse map $\Phi^{-1} : U \rightarrow \tilde{U}$. Then, $\tilde{\sigma} = \sigma \circ \Phi : \tilde{U} \rightarrow \mathbb{R}^3$ is a regular surface patch.

§4.3 Smooth Map

In this section, will define the notion of smooth map $f : S_1 \rightarrow S_2$, where S_1 and S_2 are smooth surfaces.

Definition 4.3.1: Diffeomorphisms

Smooth maps $f : S_1 \rightarrow S_2$, which are bijective and whose inverse map $f^{-1} : S_2 \rightarrow S_1$ is smooth are called diffeomorphisms.

Proposition 4.3.1

Let $f : S_1 \rightarrow S_2$ be a diffeomorphism. If σ_1 is an allowable surface patch on S_1 , then $f \circ \sigma_1$ is an allowable surface patch on S_2 .

§4.4 Tangents and derivatives

Definition 4.4.1: Tangent

A tangent vector to a surface S at a point $\mathbf{p} \in S$ is the tangent vector at \mathbf{p} of a curve in S passing through \mathbf{p} . The tangent space $T_{\mathbf{p}}S$ of S at \mathbf{p} is the set of all tangent vectors to S at \mathbf{p} .

Proposition 4.4.1

Let $\sigma : U \rightarrow \mathbb{R}^3$ be a patch of a surface S containing a point $\mathbf{p} \in S$, let (u, v) be coordinates in U . The tangent space to S at \mathbf{p} is a vector subspace of \mathbb{R}^3 spanned by vectors σ_u and σ_v (the derivatives are evaluated at the point $(u_0, v_0) \in U$ such that $\sigma(u_0, v_0) = \mathbf{p}$)

Since, by the above proposition we can see that the tangent space is 2D and will be called *tangent plane* from now on.

Remember This text only contains important theorems and definitions from the textbook **Elementary Differential Geometry**. And some of the problems for the book are also discussed (non-trivial problems).