

# Solution to HW7

①

1) Try yourself.

2) Fix  $m \neq 0$ . Then  $\lim_{y \rightarrow 0} f(my, y) = \frac{m}{1+m^2}$

Thus  $f$  is ~~not~~ not continuous at  $(0,0)$

Given  $v = (a, b)$

$$f'((0,0); v) = \lim_{t \rightarrow 0} \frac{f(tv) - f(0)}{t}$$

$$= \lim_{t \rightarrow 0} \frac{f(ta, tb)}{t} = \lim_{t \rightarrow 0} \frac{t^3 ab^2}{t(ta^2 + t^4 b^4)}$$

$$= \lim_{t \rightarrow 0} \frac{ab^2}{a^2 + tb^4} = \begin{cases} \frac{b^2}{a} & \text{if } a \neq 0 \\ 0 & \text{if } a = 0 \end{cases}$$

3) See below.

4) (i)  $f(p+tv) = f(1+t, 1+2t, 1+3t)$

$$= (1+t)^2 + (1+2t)(1+3t) + (1+t)(1+3t)^2$$

$$= 3 + 2t + 5t + 7t + \text{higher order terms}$$

$$f(p) = 3$$

$$\text{Hence, } f'(p; v) = \lim_{t \rightarrow 0} \frac{f(p+tv) - f(p)}{t} = 2 + 5 + 7 = 14$$

(ii)  $\text{grad } f = (f_x, f_y, f_z) = (y+z, x+z, x+y)$

$$\Rightarrow \text{grad } f(p) = (1, 2, 1)$$

$$\Rightarrow f'(p; v) = \text{grad } f(p) \cdot v = (1, 2, 1) \cdot (0, 1, 1) = 3$$

(iii)  $\text{grad } f = (2xy + y^2, x^2 + 2xy) \Rightarrow \text{grad } f(p) = (8, 5)$

$$\Rightarrow f'(p; v) = (8, 5) \cdot (1, 1) = 13$$

5) See below.

②

6) Recall: If  $f(x) = (f_1(x), \dots, f_m(x)) : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$  is smooth then

$$Jf = \begin{pmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \dots & \frac{\partial f_1}{\partial x_n} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \dots & \frac{\partial f_2}{\partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \frac{\partial f_m}{\partial x_2} & \dots & \frac{\partial f_m}{\partial x_n} \end{pmatrix}$$

Thus when  $n=1$ ,  $Jf = \begin{pmatrix} \frac{\partial f_1}{\partial x_1} \\ \vdots \\ \frac{\partial f_m}{\partial x_1} \end{pmatrix}$

Therefore, if  $\alpha : (-\epsilon, \epsilon) \rightarrow U \subseteq \mathbb{R}^n$  is a curve,

$\alpha(t) = (\alpha_1(t), \dots, \alpha_n(t))$  then

$$J\alpha(0) = \begin{pmatrix} \alpha_1'(0) \\ \vdots \\ \alpha_n'(0) \end{pmatrix} = \alpha'(0) \text{ written as a column vector.}$$

The rest is chain rule.


7) (i)  $f'(x) = e^x > 0$  Hence, we are done by the inverse function theorem. Here  $\text{Im}(f) = (0, \infty)$  clearly we can directly prove that  $\log : (0, \infty) \rightarrow \mathbb{R}$  is a smooth inverse of  $e^x$ .

(ii) One can directly find  $f^{-1}$  and note that  $f, f^{-1}$  are smooth. Or just find  $f'(x)$  and show that it is nonzero  $\forall x \in \mathbb{R}$  if  $f$  were to be a diffeomorphism onto image.

$$f'(x) = \frac{(x^2+1) - 2x \cdot x}{(x^2+1)^2} = \frac{1-x^2}{(x^2+1)^2}$$

Since  $f'(\pm 1) = 0$   $f$  cannot be a diffeomorphism

onto its image.  
 Ex: Check ~~if~~ if  $f$  is a homeomorphism onto its image. ③



(iii)  $f'(0) = 0$  Hence  $f$  cannot be a diffeomorphism onto image. Note:  $f^{-1}(x) = x^{1/3}$  is continuous. Hence,  $f$  is a homeomorphism  $\mathbb{R} \rightarrow \mathbb{R}$ .

$$(w) \quad Jf = \begin{pmatrix} \cos y & -x \sin y \\ \sin y & x \cos y \end{pmatrix}$$

$$\Rightarrow \det Jf = x > 0$$

Hence, by the inverse function theorem we are done.

Note: This is just transformation of polar co-ordinates into Cartesian co-ordinates.

$$8. (i) \quad \varphi_x = (1, 0, y), \quad \varphi_y = (0, 1, x)$$

$$\Rightarrow \varphi_x \times \varphi_y = \begin{vmatrix} i & j & k \\ 1 & 0 & y \\ 0 & 1 & x \end{vmatrix} = -y\vec{i} - x\vec{j} + \vec{k} \neq \vec{0} \quad \forall (x, y)$$

Hence,  $\varphi$  is an allowable surface patch.

Well one needs to check that  $\varphi$  is a homeomorphism onto its image  $S = \{(x, y, z) : z = xy\}$ . Clearly,  $\varphi$  is bijective onto  $S$  and  $\varphi^{-1}(x, y, z) = (x, y)$  which is continuous.

$$(ii) \quad \varphi_y = (0, 2y, 3y^2) \Rightarrow \varphi_y(0, 0) = 0.$$

Hence,  $\varphi$  cannot be an allowable surface patch.

Check if it is a surface patch.

(4)

(iii) Clearly  $\varphi$  is ~~not onto image~~ not even

$$1-1: \quad \varphi(0, y) = \varphi(-1, y).$$

Hence,  $\varphi$  is not a surface patch.

(iv) •  $\varphi$  is a surface patch for the cone

$z = \sqrt{x^2 + y^2}$  with vertex ~~=~~  $(0, 0, 0)$  removed.

•  $\varphi$  is smooth.

$$\varphi_x = \left(1, 0, \frac{x}{\sqrt{x^2 + y^2}}\right), \quad \varphi_y = \left(0, 1, \frac{y}{\sqrt{x^2 + y^2}}\right)$$

Clearly  $\varphi_x \times \varphi_y \neq \vec{0} \quad \forall (x, y) \in \mathbb{R}^2 \setminus \{(0, 0)\}$ .

Hence,  $\varphi$  is an allowable surface patch.

(v) •  $\varphi$  is a surface patch for the cylinder  $x^2 + y^2 = 1$   
with one line:  $x=1, y=0$  removed.

•  $\varphi$  is smooth

$$\varphi_x = (-\sin x, \cos x, 0)$$

$$\varphi_y = (0, 0, 1)$$

$$\Rightarrow \varphi_x \times \varphi_y = \sin x \vec{j} + \cos x \vec{k} \neq 0 \quad \forall (x, y)$$

$\Rightarrow \varphi$  is an allowable surface patch.

(3) and (5): The crucial thing to use is  
the Mean Value Theorem: If  $f: [a, b] \rightarrow \mathbb{R}$

is  $C^1$  then  $\frac{f(b) - f(a)}{b - a} = f'(c)$  for some

$c \in (a, b)$ .

For (3) suppose  $\{x_k\}$  is a sequence in  $\mathcal{U}$  (5)  
 converging to  $x$  where

$$x_k = (a_{1k}, a_{2k}, \dots, a_{nk}) \quad \text{and}$$

$$x = (a_1, a_2, \dots, a_n).$$


For simplicity assume  $n=2$ .

Write

$$a_{1k} = a_1 + \epsilon_k$$

$$a_{2k} = a_2 + \delta_k$$

We have  $\epsilon_k \rightarrow 0$ ,  $\delta_k \rightarrow 0$  and we want to show that  $f(a_1 + \epsilon_k, a_2 + \delta_k) \rightarrow f(a_1, a_2)$ .



Now,

$$f(a_1 + \epsilon_k, a_2 + \delta_k) - f(a_1, a_2)$$

$$= \left[ f(a_1 + \epsilon_k, a_2 + \delta_k) - f(a_1 + \epsilon_k, a_2) \right] + \left[ f(a_1 + \epsilon_k, a_2) - f(a_1, a_2) \right]$$

$$= f_y(a_1 + \epsilon_k, a_2 + \delta'_k) \cdot \delta_k + f_x(a_1 + \epsilon'_k, a_2) \cdot \epsilon_k$$

for some  $0 < \delta'_k < \delta_k$  and  $0 < \epsilon'_k < \epsilon_k$

Now, as  $f_x, f_y$  are continuous

$$f_y(a_1 + \epsilon_k, a_2 + \delta'_k) \rightarrow f_y(a_1, a_2) \quad \text{and}$$

$$f_x(a_1 + \epsilon'_k, a_2) \rightarrow f_x(a_1, a_2) \quad \text{as } \begin{matrix} \epsilon_k \rightarrow 0 \\ \delta_k \rightarrow 0 \end{matrix}$$

Hence,  $\lim_{k \rightarrow \infty} \left[ f(a_1 + \epsilon_k, a_2 + \delta_k) - f(a_1, a_2) \right] = 0$ .

[Do (5) similarly.]