

# HW1 / Hints to Answers

①

2.  $\varphi: t \mapsto (t, b, c)$  is continuous

$$g = f \circ \varphi.$$

3. (a)  $xy$ ,  $x^2 + y^2 + 1$  are continuous &  $x^2 + y^2 + 1$  is never zero.

Note: If  $f, g$  are continuous then  $f/g$  is continuous wherever  $g \neq 0$

(b)  $f$  is not continuous

If  $P_n = (x_n, mx_n) \rightarrow (0, 0)$  then

$$f(P_n) = \frac{m}{1+m^3}$$

For different sequences  $P_n \rightarrow (0, 0)$  we have different limits

(c) Let  $P_n = (x_n, y_n)$  any sequence,  $P_n \rightarrow (0, 0)$ .  
Write  $x_n = r_n \cos \theta_n$ ,  $y_n = r_n \sin \theta_n$ .  
 $\theta_n$  need not be unique, we don't care.


$$f(P_n) = r_n^2 (\cos^4 \theta_n + \sin^4 \theta_n).$$

Since  $P_n \rightarrow (0, 0)$ ,  $r_n \rightarrow 0$ . Note

$$|\cos^4 \theta_n + \sin^4 \theta_n| \leq 2. \quad \text{Thus } f(P_n) \rightarrow 0.$$

4. It is the composition of  $f$  and

$$\mathbb{R} \rightarrow \mathbb{R} \\ x \mapsto |x|$$



5. Define  $f: \mathbb{C} \rightarrow \mathbb{R} \\ (x, y) \mapsto y$

- One has check
- $f$  continuous
  - $f$  bijective
  - $f^{-1}$  continuous.

$f$  is just projection on the  $y$ -coordinate axis.

$$f^{-1}: y \mapsto (y^2, y)$$


6. Let  $f: \mathbb{R} \rightarrow [0, 1)$  be any homeomorphism

Then define  $g: \mathbb{R}^2 \rightarrow \mathbb{D} \\ (x, y) \mapsto \begin{cases} f(r)(x, y)/r & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$

where  $r = \sqrt{x^2 + y^2}$ .

For  $f$  one could take  $f(x) = \frac{e^x - 1}{e^x + 1}$

Break the circle in 4 equal parts



Map each part onto homeomorphically onto the sides of the square.

Consider  $f: [0, 1) \rightarrow S^1 = \{(x, y) : x^2 + y^2 = 1\} \\ t \mapsto (\cos 2\pi t, \sin 2\pi t)$