

2. Designate the corners of a square,  $l$  on a side, in clockwise order,  $A, B, C, D$ . Put charges  $2q$  at  $A$  and  $-3q$  at  $B$ . Determine the value of the line integral of  $\mathbf{E}$ , from point  $C$  to point  $D$ . (No actual integration needed!) What is the numerical answer if  $q = 10^{-9}C$  and  $l = 5\text{cm}$ ? [2.5]

## ASSIGNMENT 9 (SOLS.)

1. Let current flowing through outer solenoid be  $I_2$ .

Assuming field inside the solenoid to be uniform, & assuming the magnitude to be the same as that of an infinite solenoid, we have,

$$B_2 = \mu_0 N_2 I_2 = \mu_0 \left( \frac{N_2}{b_2} \right) I_2$$


$$\text{since, } n_2 = \frac{N_2}{b_2}$$

$\therefore$  Flux through inner solenoid,

$$\begin{aligned} \phi_{12} &= N_1 B_2 A_1 \quad (\text{+ve flux of inner solenoid}) \\ &= N_1 \mu_0 \frac{N_2}{b_2} I_2 \cdot \pi a_1^2 \\ &= \left( \frac{\mu_0 \pi a_1^2 N_1 N_2}{b_2} \right) I_2 \end{aligned}$$

$\therefore$  Mutual inductance ,

$$M = \frac{\mu_0 \pi a_1^2 N_1 N_2}{b_2}$$



## PHY102 : Quiz 1

1. A spherical charge distribution has a density  $\rho$  that is constant from  $r = 0$  out to  $r = R$  and is zero beyond. What is the electric field for all values of  $r$ , both less than and greater than  $R$ ? [2.5]

You may also have calculated the flux through the outer solenoid due to the inner solenoid.

Try it & see if they match!

2. Magnetic field inside solenoid,

$$B = \mu_0 n_1 I_1 = \mu_0 \frac{N_1 I_1}{l}$$

flux,  $\phi = BA = \frac{\mu_0 N_1 I_1 A}{l}$ .

$\therefore$  Self Inductance,  $L_1 = \frac{N_1 \phi}{I_1} = \frac{\mu_0 N_1^2 A}{l}$

Now,  $\phi'$  is the magnetic flux through each turn of the outer coil.

$$\phi' = B A', \quad B: \text{mag. field Due to inner solenoid.}$$

$$A': \text{area of outer coil}$$

$$= A.$$

$$\therefore \phi' = \frac{\mu_0 N_1 I_1}{l} \cdot A$$

$\therefore$  Mutual inductance,

$$M_2 = \frac{N_2 \phi'}{I_1} = \frac{\mu_0 N_1 N_2 A}{l}$$

Now, self inductance of inner solenoid,  $L_1 = \frac{\mu_0 N_1^2 A}{l}$

2. Designate the corners of a square,  $l$  on a side, in clockwise order,  $A, B, C, D$ . Put charges  $2q$  at  $A$  and  $-3q$  at  $B$ . Determine the value of the line integral of  $\mathbf{E}$ , from point  $C$  to point  $D$ . (No actual integration needed!) What is the numerical answer if  $q = 10^{-9}C$  and  $l = 5\text{cm}$ ? [2.5]

Self inductance due to <sup>outer</sup> coil,  $L_2 \approx \frac{\mu_0 N_2 A}{l}$ .

$$\therefore M \approx \sqrt{L_1 L_2}$$

(This is an idealization  $\rightarrow$  all of the magnetic flux produced by solenoid passes through the outer coil).

3.  $E_x = 0, E_y = E_0 \sin(kx + \omega t), E_z = 0$   
 $B_x = 0, B_y = 0, B_z = -\frac{E_0}{c} \sin(kx + \omega t)$

$$\vec{\nabla} \cdot \vec{E} = \frac{\partial E_x}{\partial x} + \frac{\partial E_y}{\partial y} + \frac{\partial E_z}{\partial z} = 0.$$

Similarly,  $\vec{\nabla} \cdot \vec{B} = 0$ .

$$\vec{\nabla} \times \vec{E} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ E_x & E_y & E_z \end{vmatrix} = \frac{1}{c} E_0 k \cos(kx + \omega t) \vec{k}$$

$$-\frac{\partial \vec{B}}{\partial t} = \frac{1}{c} E_0 k \cos(kx + \omega t) \vec{k}$$

To require,  $\vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{\partial t} \Rightarrow k = \frac{\omega}{c} \Rightarrow \boxed{\omega = ck}$

Use this to show that,  $\vec{\nabla} \times \vec{B} = \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t}$ .

PHY102 : Quiz 1

1. A spherical charge distribution has a density  $\rho$  that is constant from  $r = 0$  out to  $r = R$  and is zero beyond. What is the electric field for all values of  $r$ , both less than and greater than  $R$ ? [2.5]

4.  $\vec{\nabla} \cdot \vec{E} = 0.$

$$\vec{\nabla} \cdot \vec{B} = (-B_0 k \sin kx \sin ky + k B_0 \sin kx \cos ky) \sin \omega t \\ = 0.$$

$$\vec{\nabla} \cdot \vec{E} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 0 & 0 & E_0 \sin kx \cos ky \sin \omega t \end{vmatrix}$$

$$= \hat{x} \left[ \frac{\partial}{\partial y} (E_0 \sin kx \cos ky \sin \omega t) - \frac{\partial}{\partial z} (0) \right].$$

$$- \hat{y} \left[ \frac{\partial}{\partial x} (E_0 \sin kx \cos ky \sin \omega t) - \frac{\partial}{\partial z} (0) \right].$$

$$+ \hat{z} \left[ \frac{\partial}{\partial x} (0) - \frac{\partial}{\partial y} (0) \right].$$

$$= -\hat{x} E_0 k \cos kx \sin ky \cos \omega t + \hat{y} E_0 k \sin kx \cos ky \cos \omega t$$

$$-\frac{\partial \vec{B}}{\partial t} = \hat{x} B_0 \sin kx \sin ky \cos \omega t - \hat{y} B_0 \sin kx \cos ky \cos \omega t$$

Now for  $\vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}$  to hold,

$$(1) \rightarrow \boxed{E_0 k = B_0 \omega} \quad (\text{equating components}).$$

Now, let's look at the other Maxwell's eqn.

$$\vec{\nabla} \times \vec{B} = \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t}$$

2. Designate the corners of a square,  $l$  on a side, in clockwise order,  $A, B, C, D$ . Put charges  $2q$  at  $A$  and  $-3q$  at  $B$ . Determine the value of the line integral of  $\mathbf{E}$ , from point  $C$  to point  $D$ . (No actual integration needed!) What is the numerical answer if  $q = 10^{-9}C$  and  $l = 5\text{cm}$ ? [2.5]

$$\frac{\partial \vec{E}}{\partial t} = -\hat{x} \omega E_0 \cos kx \cos \theta y \sin \omega t.$$

$$\vec{\nabla} \times \vec{B} = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ B_0 \cos kx \cos \theta y \sin \omega t & -B_0 \sin kx \cos \theta y \sin \omega t & 0 \end{vmatrix}$$

$$= \hat{x} (0 - 0) + \hat{y} (0 - 0)$$

$$+ \hat{z} (-B_0 k \sin kx \cos \theta y \sin \omega t - B_0 k \cos kx \cos \theta y \sin \omega t).$$

$$= -\hat{z} 2B_0 k \cos kx \cos \theta y \sin \omega t.$$

∴ For  $\vec{\nabla} \times \vec{B} = \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t}$  to hold,

$$2B_0 k = \mu_0 \epsilon_0 \cdot \omega E_0, = \frac{\omega E_0}{c^2} \quad (\because c^2 = \frac{1}{\mu_0 \epsilon_0})$$

$$\Rightarrow \boxed{2B_0 k = \frac{\omega E_0}{c^2}} \quad \text{---(2)}$$

$$\therefore (1) \div (2) \Rightarrow \frac{E_0}{2B_0} = \frac{B_0}{E_0/c^2} \Rightarrow E_0^2 = \underline{\underline{2B_0 c^2}}$$

$$\therefore E_0 = \sqrt{2} c B_0.$$

$$\text{Substituting for } E_0 \text{ in (1)} \Rightarrow k \sqrt{2} c B_0 = B_0 \omega. \\ \therefore \omega = \sqrt{2} ck.$$

PHY102 : Quiz 1

1. A spherical charge distribution has a density  $\rho$  that is constant from  $r = 0$  out to  $r = R$  and is zero beyond. What is the electric field for all values of  $r$ , both less than and greater than  $R$ ? [2.5]

5. Proof of  $\vec{E} \cdot \vec{B}' = \vec{E}' \cdot \vec{B}$  given in text.

Basically, for a frame F' moving with speed  $v$  in the  $\hat{x}$  direction relative to F, the transformation equations are,

$$E_x' = E_x ; \quad E_y' = \gamma(E_y - vB_z) ; \quad E_z' = \gamma(E_z + vB_x)$$

$$B_x' = B_x ; \quad B_y' = \gamma(B_y + \frac{v}{c^2}E_x) ; \quad B_z' = \gamma(B_z - \frac{v}{c^2}E_y)$$

$$\vec{E}' \cdot \vec{B}' = E_x'B_x' + E_y'B_y' + E_z'B_z'$$

Using the transformations given above it is easy to see that,  $\vec{E}' \cdot \vec{B}' = \vec{E} \cdot \vec{B}$ .

To show  $E'^2 - cB'^2 = E^2 - cB^2$ , we can use the same transformations as above & do it.

Or, as suggested by Purcell in prob 9.12, we can break  $\vec{E}$  &  $\vec{B}$  into  $H^{\perp}$  &  $L^{\perp}$  vectors

$$\vec{E} = \vec{E}_{||} + \vec{E}_{\perp} \quad \& \quad \vec{B} = \vec{B}_{||} + \vec{B}_{\perp}$$

$$\therefore E'^2 - cB'^2 = \vec{E}' \cdot \vec{E}' - c \vec{B}' \cdot \vec{B}'$$

$$= (\vec{E}'_{||} + \vec{E}'_{\perp}) \cdot (\vec{E}'_{||} + \vec{E}'_{\perp})$$

$$- c (\vec{B}'_{||} + \vec{B}'_{\perp}) \cdot (\vec{B}'_{||} + \vec{B}'_{\perp})$$

2. Designate the corners of a square,  $l$  on a side, in clockwise order,  $A, B, C, D$ . Put charges  $2q$  at  $A$  and  $-3q$  at  $B$ . Determine the value of the line integral of  $\mathbf{E}$ , from point  $C$  to point  $D$ . (No actual integration needed!) What is the numerical answer if  $q = 10^{-9}C$  and  $l = 5\text{cm}$ ? [2.5]

$$\text{Now, } \vec{E}_{||}' \cdot \vec{E}_\perp' = 0, \quad \vec{B}_{||}' \cdot \vec{B}_\perp' = 0 = \vec{B}_2' \vec{B}_1' \\ = \vec{E}_\perp' \cdot \vec{E}_{||}'$$

$$\therefore E'^2 - cB'^2 = (\vec{E}_{||}' \cdot \vec{E}_{||}' + \vec{E}_\perp' \cdot \vec{E}_\perp') - c(\vec{B}_{||}' \cdot \vec{B}_{||}' + \vec{B}_\perp' \cdot \vec{B}_\perp')$$

Now, in vector form, the transformation eqs. are,

$$\vec{E}_{||}' = \vec{E}_{||} \quad ; \quad \vec{E}_\perp' = \gamma(\vec{E}_\perp + \vec{v} \times \vec{B}_\perp) \\ \vec{B}_{||}' = \vec{B}_{||} \quad ; \quad \vec{B}_\perp' = \gamma(\vec{B}_\perp - \frac{\vec{v}}{c^2} \times \vec{E}_\perp).$$

$$\therefore \vec{E}_{||}' \cdot \vec{E}_{||}' - c\vec{B}_{||}' \cdot \vec{B}_{||}' = \vec{E}_{||} \cdot \vec{E}_{||} - c\vec{B}_\perp \cdot \vec{B}_\perp$$

$$\vec{E}_\perp \cdot \vec{E}_\perp - c\vec{B}_\perp \cdot \vec{B}_\perp$$

$$= \gamma^2 (\vec{E}_\perp + \vec{v} \times \vec{B}_\perp) \cdot (\vec{E}_\perp + \vec{v} \times \vec{B}_\perp) \\ - \gamma^2 c(\vec{B}_\perp - \frac{\vec{v}}{c^2} \times \vec{E}_\perp) \cdot (\vec{B}_\perp - \frac{\vec{v}}{c^2} \times \vec{E}_\perp).$$

$$= \gamma^2 \left[ \vec{E}_\perp \cdot \vec{E}_\perp + \vec{E}_\perp \cdot (\vec{v} \times \vec{B}_\perp) + (\vec{v} \times \vec{B}_\perp) \cdot \vec{E}_\perp \right. \\ \left. + (\vec{v} \times \vec{B}_\perp) \cdot (\vec{v} \times \vec{B}_\perp) - c\vec{B}_\perp \cdot \vec{B}_\perp \right. \\ \left. - \frac{1}{c^2} \vec{B}_\perp \cdot (\vec{v} \times \vec{E}_\perp) - \frac{1}{c^2} (\vec{v} \times \vec{E}_\perp) \cdot \vec{B}_\perp \right. \\ \left. + \frac{1}{c^4} (\vec{v} \times \vec{E}_\perp) \cdot (\vec{v} \times \vec{E}_\perp) \right].$$

Now,  $\vec{E}_{||}$  is parallel to  $\vec{v}$  by definition.

PHY102 : Quiz 1

1. A spherical charge distribution has a density  $\rho$  that is constant from  $r = 0$  out to  $r = R$  and is zero beyond. What is the electric field for all values of  $r$ , both less than and greater than  $R$ ? [2.5]

$$\therefore \vec{E}_1 \text{ is } \perp \text{ to } \vec{v} \quad \therefore \vec{v} \cdot \vec{E}_1 = \vec{E}_1 \cdot \vec{v} = 0.$$

$$\therefore \vec{B}_1 \cdot (\vec{v} \times \vec{B}_1) = (\vec{v} \times \vec{B}_1) \cdot \vec{B}_1 \text{ Similarly, } \vec{v} \cdot \vec{B}_2 = \vec{B}_2 \cdot \vec{v} = 0$$

$$\therefore \underbrace{(\vec{v} \times \vec{E}_1)}_A \cdot \underbrace{(\vec{v} \times \vec{E}_2)}_B = \vec{v} \cdot [\vec{E}_1 \times (\vec{v} \times \vec{E}_2)]_C \quad (\because \vec{A} \cdot (\vec{B} \times \vec{C}) = \vec{B} \cdot (\vec{C} \times \vec{A}))$$

$$\text{Now, } \underbrace{\vec{E}_1 \times (\vec{v} \times \vec{E}_2)}_A \cdot \underbrace{\vec{v} \times \vec{E}_1}_{B-C} = \vec{v} (\vec{E}_1 \cdot \vec{E}_2) - \vec{E}_1 (\vec{v} \cdot \vec{E}_2) \quad (\because \vec{A} \times (\vec{B} \times \vec{C}) = \vec{B}(\vec{A} \cdot \vec{C}) - \vec{C}(\vec{A} \cdot \vec{B}))$$

$$= \vec{v} E_1^2$$

$$\therefore (\vec{v} \times \vec{E}_1) \cdot (\vec{v} \times \vec{E}_2) = (\vec{v} \cdot \vec{v}) E_1^2 = v^2 E_1^2.$$

$$\Rightarrow (\vec{v} \times \vec{E}_1)^2 = v^2 E_1^2.$$

$$\text{Similarly, } (\vec{v} \times \vec{B}_2) \cdot (\vec{v} \times \vec{B}_1) = v^2 B_2^2.$$

$$\therefore E_1^2 - c^2 B_2^2 = v^2 [E_1^2 + 2\vec{E}_1 \cdot (\vec{v} \times \vec{B}_2) + v^2 B_2^2 - c^2 B_2^2 + 2\vec{B}_2 \cdot (\vec{v} \times \vec{E}_1) - \frac{1}{c^2} \cdot v^2 E_1^2].$$

$$\text{But, } \vec{E}_1 \cdot (\vec{v} \times \vec{B}_2) = -\vec{B}_2 \cdot (\vec{v} \times \vec{E}_1) \quad (\because \vec{A} \cdot (\vec{B} \times \vec{C}) = -\vec{C} \cdot (\vec{B} \times \vec{A}))$$

$$\therefore E_1^2 - c^2 B_2^2 = v^2 [E_1^2 (1 - \frac{v^2}{c^2}) + v^2 (B_2^2 - \frac{v^2}{c^2} E_1^2) - c^2 B_2^2 (1 - \frac{v^2}{c^2})]$$

2. Designate the corners of a square,  $l$  on a side, in clockwise order,  $A, B, C, D$ . Put charges  $2q$  at  $A$  and  $-3q$  at  $B$ . Determine the value of the line integral of  $\mathbf{E}$ , from point  $C$  to point  $D$ . (No actual integration needed!) What is the numerical answer if  $q = 10^{-9} C$  and  $l = 5\text{cm}$ ? [2.5]

$$\therefore \bar{E}_1' - c^2 \bar{B}_2' = \gamma^2 \left[ \frac{\bar{E}_1^2}{\gamma^2} - c^2 \frac{\bar{B}_2^2}{\gamma^2} \right] = \bar{E}_1^2 - c^2 \bar{B}_2^2$$

$$\therefore \bar{E}^2 - c^2 \bar{B}^2 = \bar{E}_1^2 - c^2 \bar{B}_2^2. \text{ Hence proved.}$$

6.  $P = 60 \text{ watt}, V = 120 \text{ volts.}$

$$\therefore \text{Current, } I = \frac{P}{V} = \frac{60}{120} = 0.5 \text{ A.}$$

$$\text{Resistance of filament, } R = \frac{V}{I} = \frac{120}{0.5} = 240 \Omega.$$

To have same  $I$  when bulb is connected in series with an inductance  $L$ , we have,

$$\text{impedance, } Z = \sqrt{R^2 + (\omega L)^2}$$

$$\therefore \text{Current, } I = \frac{V'}{\sqrt{R^2 + (\omega L)^2}} \quad \text{with } V' = 240 \text{ volt}$$

$$\Rightarrow 0.5 = \frac{240}{\sqrt{240^2 + (\omega L)^2}} \Rightarrow (\omega L)^2 = (480)^2 - (410)^2$$


$$\Rightarrow \omega L \approx 415.7 \Omega \quad \omega = 2\pi f = 2\pi(60)$$

$$\therefore L \approx \frac{415.7}{2\pi \cdot 60} \approx 1.10 \text{ henry.}$$

PHY102: Quiz 1

1. A spherical charge distribution has a density  $\rho$  that is constant from  $r = 0$  out to  $r = R$  and is zero beyond. What is the electric field for all values of  $r$ , both less than and greater than  $R$ ? [2.5]

7. Let's do the serial RLC chg the Purcell way.



$V$  is +ve if upper capacitor plate is truly charged. +ve current direction defined by the arrow. Then,

$$I := -\frac{d\phi}{dt} \quad ; \quad \phi : \mathcal{C}V \quad ; \quad V = \int \frac{d\phi}{dt} + PR$$

$$I = -\frac{d}{dt}(cv) = -c \frac{dv}{dt}$$

$$\therefore V = -LC \cdot \frac{d^2V}{dt^2} = CR \cdot \frac{dV}{dt}$$

$$\Rightarrow \frac{d^2V}{dt^2} + \frac{R}{L} \frac{dV}{dt} + \left(\frac{1}{LC}\right)V = 0. \quad (1)$$

we can proceed to solve this ala Purcell (Section)

Now consider the parallel LCR circuit.



We still have,  $\theta_{2\text{C}}$

$$4 \quad I_1 = -\frac{d\phi}{dt} = -c \frac{dV}{dt}$$

Now,  $V = R'(I_1 + I_2)$  Also,  $V = -L \frac{dI_2}{dt}$ . (Note the sign.)

~~2. Weißer Hirsch~~

2. Designate the corners of a square,  $l$  on a side, in clockwise order,  $A, B, C, D$ . Put charges  $2q$  at  $A$  and  $-3q$  at  $B$ . Determine the value of the line integral of  $\mathbf{E}$ , from point  $C$  to point  $D$ . (No actual integration needed!) What is the numerical answer if  $q = 10^{-9}C$  and  $l = 5\text{cm}$ ? [2.5]

$$\begin{aligned}\therefore \frac{dV}{dt} &= R' \left( \frac{dI_1}{dt} + \frac{dI_2}{dt} \right) \\ &= R' \left[ -C \frac{d^2V}{dt^2} - \frac{V}{L} \right].\end{aligned}$$

$$\Rightarrow \frac{dV}{dt} = -R'C \frac{d^2V}{dt^2} - \frac{R'V}{L}.$$

$$\Rightarrow \frac{d^2V}{dt^2} + \frac{1}{RC} \frac{dV}{dt} + \frac{V}{LC} = 0. \quad \text{---(2)}$$

Comparing eq. (1) & (2), they are same

$$\text{If, } \frac{R}{L} = \frac{1}{RC} \Rightarrow \boxed{R' = \frac{L}{RC}}.$$