


Tutorial # 4

(a) $\mu = \frac{0.01 \text{ Kg}}{2.5 \text{ m}} ; T = 10 \text{ N}$


$$v = \sqrt{\frac{T}{\mu}} = \sqrt{\frac{10 \times 2.5}{0.01}} = \sqrt{2500} \text{ m/s}$$

freq. $\boxed{v_0 = \frac{v}{\lambda}}$ for fundamental mode $\frac{\lambda}{2} = L$

$$v_0 = \frac{v}{2L} = \frac{\sqrt{2500}}{2 \times 2.5} = \sqrt{\frac{2500}{25}} = 10 \text{ Hz}$$



(b)



$$\Rightarrow \text{Next node } \left(\frac{\lambda_n}{2}\right) = \frac{L}{n}$$

$$\boxed{\lambda_n = \frac{2L}{n}}$$

$$\Rightarrow \text{Surviving } \lambda_n; n = 5, 10, 15, \dots \infty$$

$$v_n = 50, 100, 150, \dots \text{ Hz}$$

Q-2

μ, T



At $t=0$ $y(x,0) = 2 \sin\left(\frac{2\pi x}{L}\right) + 3 \sin\left(\frac{\pi x}{L}\right)$

(a) At $t=0$ all energy is potential (deformation of string)


Potential energy density $\frac{dU}{dx} \approx \frac{1}{2} T \left(\frac{\partial y}{\partial x}\right)^2$

- Total energy stored in (energy conservation)

$$\boxed{E_{\text{tot}} = U(t=0) = \frac{1}{2} T \int_0^L \left(\frac{\partial y}{\partial x}\right)^2 dx}$$

$$\left(\frac{\partial y}{\partial x}\right)^2 = \left[\frac{4\pi}{L} \cos\left(\frac{2\pi x}{L}\right) + \frac{3\pi}{L} \cos\left(\frac{\pi x}{L}\right)\right]^2$$

\Rightarrow Cross terms in $\left(\frac{\partial y}{\partial x}\right)^2$ will NOT contribute to integral \int_0^L



$$E_{tot} = \frac{1}{2} T \frac{\pi^2}{L^2} \left[16 \int_0^L \cos^2\left(\frac{2\pi x}{L}\right) dx + 9 \int_0^L \cos^2\left(\frac{\pi x}{L}\right) dx \right]$$



$$E_{tot} = \frac{25 \pi^2 T}{4L}$$

(b) Superposition of two-modes: two-standing waves

$$y(x,t) = 3 \sin\left(\frac{\pi x}{L}\right) \cos(\omega_1 t) + 2 \sin\left(\frac{2\pi x}{L}\right) \cos(\omega_2 t)$$

~~$$\omega = kv \Rightarrow \omega = \frac{2\pi v}{\lambda}$$~~


$$v = \frac{v}{\lambda} \Rightarrow \omega = \frac{2\pi v}{\lambda} \quad ; \quad v = \sqrt{\frac{T}{\mu}}$$



$$\omega_1 = \frac{2\pi v}{\lambda_1}$$

$$+ \quad \omega_2 = \frac{2\pi v}{\lambda_2}$$

Two standing waves:



+



$$\mu = 0.1 \text{ g/cm} = 0.01 \text{ kg/m}$$

$$T = 400 \text{ Hz}$$

$$\text{wave speed } v = \sqrt{\frac{T}{\mu}}$$

$$\text{Amplitude } A = 0.01 \text{ m}$$

$$\omega = 100 \text{ Hz}$$

$$\begin{aligned} \text{Time average energy flux } \langle P(t) \rangle &= 2\pi^2 A^2 \mu \omega^2 v \text{ J/s} \\ &= \frac{1}{2} (\mu \omega^2 A^2) v \end{aligned}$$

$$\langle P(t) \rangle = 39.48 \text{ W}$$

